

The Incredible Gastrointestinal Barrier System and Permeability: Laboratory Assessment and Insights for Intervention

by David Quig, PhD

References

1. Moran AP, et al. Sweet-talk: role of host glycosylation in bacterial pathogenesis of the gastrointestinal tract. *Gut*. 2011;60:1412-25.
2. Bischoff SC, et al. Intestinal permeability- a new target for disease prevention and therapy. *BMC Gastroenterol*. 2014;14:189.
3. Pendyala S. A high-fat diet is associated with endotoxemia that originates from the gut. *Gastroenterol*. 2012;142:1100-01.
4. Yu Lch, et al. Host-microbial interactions and regulation of intestinal epithelial barrier function: from physiology to pathology. *World J Gastroint Physiol*. 2012;15:27-43.
5. Thangaraju M, et al. Coupled receptor for the bacterial fermentation product butyrate and functions as a tumor suppressor in colon. *Cancer Research*. 2009;69:7 DOI: 10.1158/008-5472.CAN-08-4466.
6. Ouwekerk J, et al. Glycobiome: bacteria and mucus epithelial interface. *Best Prac Res. Clin Gastroenterol*. 2013;27:25-38.
7. Ploger S, et al. Microbial butyrate and its role for barrier function in the gastrointestinal tract. *Ann NY Acad Sci*. 2012;1258:52-9.
8. Matt S, et al. Butyrate and dietary soluble fiber improve neuroinflammation associated with aging in mice. *Front Immunol*. 2018;19 DOI: 10.3389/fimmu.2018.01832.
9. Lopetuso LR, et al. Commensal clostridia: leading players in the maintenance of gut homeostasis. *Gut Pathogens* 2013;5:23
10. Chen HM, et al. Decreased dietary fiber intake and structural alterations of gut microbiota in patients with advanced colorectal adenoma. *Am J Clin Nutr*. 2013;97:1044-52.
11. Jost T, et al. Vertical mother-neonate transfer of maternal gut bacteria via breastfeeding. *Environ Microbiol*. 2013;8.
12. Tenover FC, et al. Antimicrobial resistant strains of *Clostridium difficile* in North America. *Antimicrob Agents Chemother*. 2012;56:2929-32.
13. Johansson ME, et al. Bacteria penetrate the normally im-penetrable inner colon mucus layer in both murine colitis models and patients with ulcerative colitis. *Gut*. 2014;63:281-91.
14. Serino M, et al. Metabolic adaptation to a high-fat diet is associated with a change in the gut microbiota. *Gut*. 2012;61:543-53.
15. Corthesy B. Multi-faceted functions of secretory IgA at mucosal surfaces. *Front Immunol* 2013;4
16. Mathia A, et al. Role of secretory IgA in the mucosal sensing of commensal bacteria. *Gut Microbes* 2014;5:688-95.
17. Kaptustino OA, et al. Resistance of the fungi *Candida* to human innate immunity. *Gig Sanit*. 2012;4:27-9.
18. Compos-Rodriguez R, et al. Stress modulates immunoglobulin A. *Front Integrat Neurosci* 2013;7
19. Andrade ME, et al. The role of immunomodulators on intestinal barrier homeostasis. *Clin Nutr*. 2015;34:1080-7.
20. Ami-Romach E, et al. Bacterial population and innate immunity-related genes in rat gastrointestinal tract are altered by vitamin A-deficient diet. *J Nutr Biochem*. 2009;20:70-7.
21. Fasano A. Zonulin and its regulation of intestinal barrier function: the biological door to inflammation, autoimmunity, and cancer. *Physiol Rev*. 2011;91:151-75.
22. Lammers KM, et al. Gliadin induces an increase in intestinal permeability and zonulin release by binding to the chemokine receptor CXCR3. *Gastroenterol*. 2008;135:194-204e.
23. Fasano A. Zonulin, regulation of tight junctions, and autoimmune diseases. *Ann NY Acad Sci*. 2012;1258:25-33.
24. Sapone A, et al. Zonulin upregulation is associated with increased gut permeability in subjects with type I diabetes and their relatives. *Diabetes* 2006;55:1443-9.
25. Pacifico L, et al. Increased circulating zonulin in children with biopsy-proven nonalcoholic fatty liver disease. *World J Gastroenterol*. 2014;20:17107-114.
26. Asmar RE, et al. Host-dependent zonulin secretion causes the impairment of small intestine barrier function after bacterial exposure. *Gastroenterol*. 2002;123:1607-15.
27. Fasano A, et al. Nonceliac gluten sensitivity. *Gastroenterol*. 2015;148:1195-204.
28. Zak-Golab, et al. Gut microbiota, microinflammation, metabolic profile, and zonulin concentration in obese and normal weight subjects. *Int J Endocrinol*. 2013.
29. Moreno-Navarrete, JM et al. Circulating zonulin, a marker of intestinal permeability, is increased in association with obesity-associated insulin resistance. *Plos One*. 2012;7:e37160
30. Lerner A, Matthias T. Changes in intestinal tight junction permeability associated with industrial food additives explain the rising incidence of autoimmune disease. *Autoimmun Rev*. 2015;14:479-89.
31. Russo F, et al. Inulin-enriched pasta improves intestinal permeability and modifies the circulating levels of zonulin and glucagon-like peptide in healthy young volunteers. *Nutr Res*. 2012;32:940-46.
32. Elamin E, et al. Ethanol impairs intestinal barrier function in humans through mitogen activated protein kinase signaling: a combined *in vivo* and *in vitro* approach. *Plos One*. DOI: 10.1371/journal.pone.0107421.